



MASL Family Plant Nutrition

Crop-Specific Series

Formulated with targeted nutrients—B Vitamins, Glutathione, Phycocyanin, Polyglutamic Acid, Seaweed Extracts, and other beneficial organic ingredients—to safeguard crop health and maximize productivity.

摩多生物科技有限公司

Shenyang MASL Biotechnology Co., Ltd



Crop-Specific Series

Field Crop Formula

Leafy Vegetable Formula

Fruit Tree Formula

Root and Tuber Formula

Fruiting Vegetable Formula



Shenyang MASL Biotechnology Co., Ltd



MASL Family Plant Nutrition Field Crop Formula



Shenyang MASL Biotechnology Co., Ltd

Field Crop Formula

MASL[®]
MOS CONCENTRAMOS EN ECO
— 摩多 —



Shenyang MASL Biotechnology Co., Ltd

Field Crop Formula



Soybean Fill Soybean Late Stage Specialty Fertilizer

Technical Specification:

Free Amino Acids $\geq 100\text{g/L}$, Ca+Mg $\geq 30\text{g/L}$

Free Amino Acids:	100g/L	Mg:	30g/L	Zn:	2.5g/L
Water-soluble Boron:	2.5g/L	Vitamin B3:	0.5g/L	/	/

Benefits:

Specifically developed for mid-to-late soybean growth needs. Uses highly active L- α free amino acids as the core, scientifically compounded with chelated Magnesium, Zinc, and Boron, precisely tailored for the flowering/pod setting to grain filling stages. Highly active L- α free amino acids promote flower bud differentiation and pod setting stability, reduce flower and pod drop, synergize with chelated Magnesium to enhance photosynthetic efficiency, providing continuous energy for grain filling. Chelated Zinc activates protein synthase activity, working with Boron to ensure pollen vitality and fertilization quality, reducing blighted grain rate, promoting efficient oil and protein accumulation, achieving full pods and increased hundred-grain weight. Specially added Vitamin B3 improves photosynthetic efficiency and energy metabolism, delays functional leaf senescence, enhances plant stress resistance, and maintains key enzyme activity. Promotes directional nutrient transport to pods, enhancing yield and quality for increased profitability.

Applicable Crops: soybean at late stage

Application Instructions:

Foliar Spray: Dilute 800-1000 times. Apply at 7-10 day intervals, for 2-3 consecutive applications.

Fertigation/Drip Irrigation: 15-30L per hectare

Cautions:

1. Avoid mixing with strongly acidic or alkaline fertilizers and pesticides, as this may affect product efficacy. If mixing with other products is intended, conduct a small-scale compatibility test first.
2. Reapplication at a reduced rate may be necessary if rainfall occurs within 4 hours of spraying.

Storage:

1. Store and transport in the original container. Keep sealed in a cool, dry place.
2. Store separately from food, drink, and feed. Keep out of reach of children and livestock.

Production Date and Lot No.: See bottle cap. Shelf Life: 36 months.

Shenyang MASL Biotechnology Co., Ltd