



MASL Family Plant Nutrition

Crop-Specific Series

Formulated with targeted nutrients—B Vitamins, Glutathione, Phycocyanin, Polyglutamic Acid, Seaweed Extracts, and other beneficial organic ingredients—to safeguard crop health and maximize productivity.

摩多生物科技有限公司

Shenyang MASL Biotechnology Co., Ltd



Crop-Specific Series

Field Crop Formula

Leafy Vegetable Formula

Fruit Tree Formula

Root and Tuber Formula

Fruiting Vegetable Formula



Shenyang MASL Biotechnology Co., Ltd



MASL Family Plant Nutrition Field Crop Formula



Shenyang MASL Biotechnology Co., Ltd

Field Crop Formula

MASL[®]
MOS CONCENTRAMOS EN ECO
— 摩多 —



Shenyang MASL Biotechnology Co., Ltd

Field Crop Formula



FieldMax Boost

Field Crops Late Stage Specialty Fertilizer

Technical Specification:

Amino Acids $\geq 100\text{g/L}$, Ca+Mg $\geq 30\text{g/L}$

Free Amino Acids:	100g/L	Mg:	30g/L	Zn	2.5g/L
Water-soluble Boron:	2.5g/L	Vitamin B5:	1g/L	/	/

Benefits:

Specifically developed for the mid-to-late growth needs of field crops. Uses highly active L- α free amino acids as the core, scientifically compounded with chelated Magnesium, Zinc, and Boron. Significantly promotes growth during the booting and grain filling stages of cereal crops (e.g., rice, wheat, corn), leading to uniform ear emergence, high panicle formation rate, full grains, reduced spikelet degeneration and empty grains, promoting starch and protein synthesis, increasing yield. Also suitable for legume crops (e.g., soybean, pea, mung bean) from pod setting to grain filling stages, helping increase pod set, reduce empty pods, and improve grain plumpness. Specially added Vitamin B5 regulates late-stage metabolic functions, delays plant senescence, and promotes efficient nutrient transfer to fruits or grains, aiding dry matter accumulation. Enhances yield and quality for increased profitability.

Applicable crops: field crops at late stage

Application Instructions:

Foliar Spray: Dilute 800-1000 times. Apply at 7-10 day intervals, for 2-3 consecutive applications.

Fertigation/Drip Irrigation: 15-30L per hectare

Cautions:

1. Avoid mixing with strongly acidic or alkaline fertilizers and pesticides, as this may affect product efficacy. If mixing with other products is intended, conduct a small-scale compatibility test first.
2. Reapplication at a reduced rate may be necessary if rainfall occurs within 4 hours of spraying.

Storage:

1. Store and transport in the original container. Keep sealed in a cool, dry place.
2. Store separately from food, drink, and feed. Keep out of reach of children and livestock.

Production Date and Lot No.: See bottle cap. Shelf Life: 36 months.